

RECOMMENDED WORK PERMISSIONS AND RESTRICTIONS MANAGEMENT AS DETERMINED BY EXPOSURE RISK AND IMPACT ON SAFE SERVICE DELIVERY

If symptomatic, symptoms develop: Isolate for 7 days since last contact with COVID-19 case. Follow public health orders of jurisdiction. If you have symptoms at Day 6, you must stay at home until symptoms are gone. If you test positive D6, quarantine for a further 7 days.

Positive COVID-19 test result – case management: Isolate 7 days, RAT self-test D6. If negative and asymptomatic RTW with additional requirements. Remain vigilant for symptoms.

	LOW-RISK CONTACTS	HIGH-RISK CONTACTS
<p>NORMAL ESSENTIAL SERVICE SETTINGS</p>	<p>Continue to work, remain vigilant for symptoms, test and isolate immediately if these occur.</p> <p>Surveillance testing:</p> <ul style="list-style-type: none"> Surveillance testing can be a useful tool in the context of a specific industry. Industry can manage their WHS risk through RAT use if deemed appropriate. <p>Additional:</p> <ul style="list-style-type: none"> Surgical masks and in other circumstances^ other PPE may be required. Break areas separated, well ventilated and Team Members (TM) eat distanced by 1.5m. Wear a mask while travelling to work. 	<p>Immediately quarantine for 7 days since last contact with COVID-19 case.</p> <ul style="list-style-type: none"> RAT self-test Day 1 and Day 6 Return to work (RTW) after 7 days of quarantine and when day 6 test result returns negative and asymptomatic. Continue to monitor for symptoms for a total 14 days from exposure Apply additional requirements on RTW as below.
<p>HIGH IMPACT ON ESSENTIAL SERVICE SETTINGS</p> <p>*Critical risk to service delivery requires senior management determination.</p> <p>Essential services/supply are disrupted.</p>	<p>Suggested controls (controls should be based on local risk assessment):</p> <ul style="list-style-type: none"> Work in surgical mask and in other circumstances^ other PPE may be required. Break areas separated# (consider cohorting of staff), well ventilated and physical distancing observed. Limit work to a single site and area (where the workplace has distinct areas). If possible, travel alone to work. If not possible, wear a mask while car-pooling or using public transport. 	<p>Asymptomatic:</p> <ul style="list-style-type: none"> Continue to work if negative day 1 RAT*. RAT at every second day until Day 6. Monitor for symptoms, test, and isolate immediately if symptoms develop. Must quarantine outside of work until cleared or negative test day 6-7. Continue to monitor for symptoms for a total 14 days from exposure. <p>If negative:</p> <ul style="list-style-type: none"> RAT self-test Day 1 and every second day until Day 6. Continue to monitor for symptoms for a total 14 days from exposure. <p>If positive RAT:</p> <ul style="list-style-type: none"> Isolate and follow protocol in red box. <p>If become symptomatic:</p> <ul style="list-style-type: none"> Immediately quarantine and take RAT test. Follow protocol in red box. <p>See additional suggested controls in low risk contacts but controls should be based on local risk assessment.</p>

^ Where fitted PPE is warranted, available and training has occurred.

* If required testing unavailable, worker must not attend the workplace.

The provider must provide an adequate place for workers to observe their breaks.